


CT CHEST Without Contrast (Low dose lung screening-LDCT)



Indications	Patient age 50-80 (private insurance), Patient age 50-77 (Medicare), greater than 20 pack year history of smoking, current smoker or quit within last 15 years, no acute signs / symptoms of lung cancer.		
	Cigar & non-cigarette smoking does NOT qualify.		
Patient Position	Supine		
Contrast	NONE		
Injection rate	NONE		
Respiration	Breath- hold		
Scan Parameters	just above lung apices through adrenals		
Special note	Preferred slice thickness & interval is listed below. Minimum 3/3mm scan & 1/1 mm reformats for scanner limitations.		

PLANE	ALGORITHM	THICKNESS	INTERVAL	DFOV	REFORMATS
Axial	Mediastinum	1.25 mm	1.25 mm	pt largest + 4 cm	
Axial	Lung	1.25 mm	1.25 mm		Х
Sagittal	Mediastinum	0.625 mm	0.625 mm		Х
Coronal	Mediastinum	0.625 mm	0.625 mm		Х
Axial	MIP	8.0 mm	3.0 mm		Х



- · LMP on pts of child-bearing age
- Shield children when possible

Any deviation from protocol MUST be Radiologist approved. Rad Director will be notified if this occurs without prior approval.